

# THE IMPACT OF MULTIPLE CONTAMINANTS ON BIOMASS DYNAMICS

## INTRODUCTION

### THE CHALLENGE

Current approaches to **environmental risk assessment (ERA)** are often limited to assessing the effect of single contaminants on single species, overlooking the effects that can occur at higher ecological scales.

Can we develop tools for ERA that allow us to evaluate how **multiple contaminants** acting on **multiple traits** among species in communities impact **biodiversity, stability and ecosystem function**?

**HANA MAYALL**

hbmayall1@sheffield.ac.uk

**SUPERVISORS: ANDREW BECKERMAN, DYLAN CHILDS, LORRAINE MALTY, CLAUDIA RIVETTI**


## METHODS

### IN SILICO EXPERIMENTAL DESIGN


- Use differential equation **food web model** to simulate dynamics of plausible **tri-trophic** food chain
- Specify inhibitory **contaminant effects** on populations via linear reductions or increases of trait rates
- Generate 2-contaminant scenarios with
  - 1 **herbicide** targeting growth and
  - 1 **pesticide** targeting either metabolism or foraging
- Measure **community biomass**
- Classify interactions** by calculating deviation from additivity of community biomass<sup>[3]</sup>

### FOOD WEB MODELLING

Biomass dynamics in a community are determined by species traits <sup>[1]</sup>




### CONTAMINANT EFFECTS




### 3 SPECIES CASE STUDY

#### VARYING TARGET TRAIT



#### MEASURING BIOMASS




Is the initial dynamics important here? If so maybe you can reduce the max on the x-axis?

### CLASSIFYING INTERACTIONS

Using Tekin *et al.*'s framework for measuring **ecological stressor interactions** <sup>[2]</sup>, which incorporates;

- Standardisation** of effect sizes (rescaling)
- Categorisation** of interaction types
- Measures **Deviation from Additivity (DA)**
  - Synergy, Buffering and Suppression



- Synergy** – joint effect greater than sum of its parts
- Additive** – no interaction
- Antagonistic buffering** – joint effect smaller than sum of its parts
- Antagonistic suppression** – effect of one contaminant masks that of the other
- Out of bounds** – joint effect less than that of each individual contaminant
- Inconclusive** – normalisation factor = 0


## CONCLUSIONS

- ★ **Community biomass** is differentially impacted depending on the **trait and trophic level** targeted by the pesticide
- ★ **Additivity, antagonistic buffering and antagonistic suppression** are the most commonly observed interaction types, with **no synergy** observed
- ★ **Antagonistic suppression** is observed when pesticides target **foraging** in the **intermediate consumer**, due to compensatory biomass dynamics inherent to the model resulting in reallocation of biomass among trophic levels

## RESULTS

### VARIATION IN...

### RESCALED DEVIATION FROM ADDITIVITY OF COMMUNITY BIOMASS



## FUTURE WORK

- Explore buffering and suppression of total biomass by exploring biomass re-allocation across trophic levels
- Apply method to stability
- Expand community complexity

## TOWARDS REDEFINING ERA

How do multiple contaminants impact biodiversity, stability and ecosystem function through...

- Target trait (*mode of action*)
- Target species (*contaminant specificity*)
- Target trophic level
- Community size, structure and complexity
- Environmental conditions

## REFERENCES

- <sup>[1]</sup> Williams, Brose & Martinez (2007) Homage to Yodzis and Innes 1992: Scaling up feeding-based population dynamics to complex ecological networks  
<sup>[2]</sup> Tekin *et al.* (2020) Using a newly introduced framework to measure ecological stressor interactions